

### Drug Facts

#### Active ingredient

Benzocaine 10% . . . . . **Purpose**  
Oral pain reliever

**Uses** temporarily relieves pain caused by: ■ canker sores ■ minor mouth sores and gum irritations ■ denture and brace pain

**Warnings** **Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in our care develops: pale, gray, or blue colored skin (cyanosis); headache; rapid heart rate; shortness of breath; dizziness or lightheadedness; fatigue or lack of energy.

**Flammable:** ■ keep away from fire or flame

**Allergy Alert:** ■ do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other “caine” anesthetics

**Stop use and ask a physician if** ■ sore mouth symptoms do not improve in 7 days ■ swelling, rash or fever develops ■ irritation, pain or redness persists or worsens

■ apply only to affected area ■ do not exceed recommended dosage ■ avoid contact with the eyes ■ do not use for more than 7 days unless directed by a physician or dentist

**Keep out of reach of children.** If swallowed, get medical help or contact a poison control center immediately.

**Do not use** ■ for teething ■ in children under 3 years of age

**Directions** remove cap and cut open tip of tube on scoremark

adults and children 3 years and older	Dry affected area. Apply a thin coat of gel with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.
children under 12 years	adult supervision should be given in the use of this product
children under 3 years	do not use

**Other information** ■ do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin-B Long Lasting Mouth Sore Gel to film, and immediately wipe the area with a moist gauze pad or tissue. ■ oxidation may darken the gel's appearance over time, but its effectiveness is unchanged  
■ contains alcohol 65% by volume ■ store at 15-30°C (59-86°F)

**Inactive ingredients** boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 38-B, tannic acid