

### Drug Facts

#### Active ingredient

Benzyl alcohol 10% . . . . . Cold sore/fever blister treatment, Oral pain reliever

#### Purpose

**Uses** temporarily relieves pain caused by: ■ cold sores/ fever blisters ■ canker sores, mouth sores ■ gum irritations

**Warnings** **Flammable:** ■ keep away from fire or flame

**Stop use and ask a physician if** ■ sore mouth symptoms do not improve in 7 days ■ condition worsens or symptoms clear up and occur again within a few days ■ swelling, rash or fever develops ■ irritation, pain or redness persists or worsens

■ apply only to affected area ■ do not exceed recommended dosage ■ avoid contact with the eyes ■ do not use for more than 7 days unless directed by a physician or dentist

**Keep out of reach of children.** If swallowed, get medical help or contact a poison control center immediately.

**Directions** remove cap and cut open tip of tube on scoremark

adults and children  
2 years and older

dry affected area. Apply with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.

children under 12 years

adult supervision should be given in the use of this product

children under 2 years

do not use, consult a physician or dentist

#### Other information

■ do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin to film, and immediately wipe the area with a moist gauze pad or tissue. ■ contains alcohol 68% by volume ■ store at 15-30°C (59-86°F)

#### Inactive ingredients

boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 37, tannic acid